Adapted from WA DOH Form 347-102

## Fax Page 1 To:

# Whitman County Health Department (509) 397-6239 (Confidential FAX line)



CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION CASE REPORT
Report STIs within 3 work days (WAC 246-101-101/301)

PATIENT INFORMATION							
LAST NAME FIRST N	AME	MIDDLE NAME DATE OF BIRTH					
		MO DAY YR					
ADDRESS ( $\square$ Unhoused or unstably housed in the p	ast 3 months) CITY	STATE ZIP CODE					
TELEPHONE EMAIL	ENGLISH SPEAKING	? ☐ Yes ☐ No *instructions on pg. 3 DIAGNOSIS DATE					
( )	Preferred Language						
SEX ASSIGNED GENDER IDENTITY	ETHNICITY	RACE CATEGORY (check all that apply)*: *Instructions on page 3  ☐ White ☐ Asian					
AT BIRTH	Wate   Indispendent Will   Inspande of Edition of A						
☐ Male     ☐ Female     ☐ Transgender FTM       ☐ Female     ☐ Nonbinary /     ☐ Other	☐ Non-Hispanic ☐ Unknown	☐ Black ☐ Other ☐ Unknown					
☐ Intersex ☐ Genderqueer ☐ Refused	Refused	☐ Native Hawaiian / Other Pacific Islander ☐ Refused					
Refused		EXTENDED RACE CODE(S)*: R R R R					
l l	DER OF SEX PARTNERS (check						
PREGNANT? ☐ Exposed to Infection ☐ N	_ 0						
	emale	nder FTM					
	Ionbinary / □ Other Senderqueer □ Unknow	1 <u> </u>					
DIAGNOSIS - DISEASE		, , ,					
GONORRHEA (lab confirmed)		SYPHILIS					
DIAGNOSIS (check one) SITES (all that apply):	TREATMENT (check all pres	cribed): STAGE (check one):					
☐ Asymptomatic ☐ Cervix	☐ Ceftriaxone: ☐ 250 mg	· - ·   - · · · · · · · · · · · · · · ·					
☐ Symptomatic, Uncomplicated ☐ Urethra ☐ Pelvic Inflammatory Disease ☐ Urine		☐ 800 mg ☐ Secondary (Rash, etc.) ☐ Early Latent (< 1 year)					
☐ Ophthalmia	☐ Azithromycin: ☐ 1 g ☐ Doxycycline: ☐ 100 mg						
☐ Disseminated ☐ Pharynx	Gentamicin: 240 mg						
☐ Other Complications: ☐ Vagina	☐ Gemifloxacin: ☐ 320 mg						
Ocular	Other:	Neurologic ☐ Otic ☐ Ocular ☐ Tertiary					
Date lested.	Date Prescribed:						
CHLAMYDIA (lab confirmed)	TDEATRAERIT (sheek all muse	TREATMENT (check one):					
DIAGNOSIS (check one)   SITES (all that apply):	:	Cribed):					
☐ Asymptomatic ☐ Cervix ☐ Symptomatic, Uncomplicated ☐ Urethra	☐ Azithromycin: ☐ 1 g☐ Doxycycline: ☐ 100 mg						
☐ Pelvic Inflammatory Disease ¦☐ Urine	Levofloxacin: 500 mg						
☐ Ophthalmia ☐ Rectum ☐ Other Complications: ☐ Phanny	Other:	Benzathine					
Uther Complications: Pharynx	i !	PCN-G: 50,000 units/kg IM x 3					
Ocular	i	Aqueous 18-24 MU/day IV					
Date Tested: Other:	Date Prescribed:	Crystalline for 10-14 days					
HERPES SIMPLEX	OTHER DISEASES	Penicillin G:					
DIAGNOSIS LABORATORY CONFIRM	L Chanciola	Other:					
☐ Genital (initial infection only) ☐ Yes ☐ Neonatal ☐ No	☐ Granuloma Inguir☐ Lymphogranulom						
PARTNER TREATMENT PLAN (check one or more o							
Providers should manage partner treatment by either treating partners in-person or by prescribing medication for patients to give to their sex partners (see side 2							
for additional information).  Turn over for Partner Treatment Plan Instructions							
in-person evaluation - Number of partners treated following medical evaluation:							
☐ Patient-delivered treatment* - Number of partners for whom provider prescribed or provided expedited partner therapy (EPT) medication pack to be delivered by the patient to their partner(s): *Patient-delivered treatment is not recommended for men who have sex with men or patients with syphilis							
REPORTING CLINIC INFORMATION							
DATE FACILITY NAME	DIAGN	OSING CLINICIAN					
•	L CITY	CTATE 7ID					
ADDRESS	CITY	STATE ZIP					

#### PARTNER MANAGEMENT PLAN INSTRUCTIONS

## **Gonorrhea or Chlamydia Infection: Partner Treatment**

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Whitman County Health Department may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Whitman County Health Department recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Whitman County Health Department: (509) 397-6280.

#### Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

## RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS\*

## **GONORRHEA -- Uncomplicated**

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs) $^{\dagger}$ 

## Alternatives for uncomplicated infections of the cervix, urethra, or rectum:

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose  $\bf OR$  Cefixime 800 mg orally as a single dose  $^{\dagger}$ 

- † If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.
- <sup>‡</sup> Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

#### **CHLAMYDIA** -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

#### Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days OR

Ethylsuccinate 800 mg PO QID for 7 days OR

Ofloxacin 300 mg PO BID for 7 days OR

Levofloxacin 500 mg PO for 7 days

#### SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

## **SYPHILIS -- LATE OR UNKNOWN DURATION**

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

DOH 347-102, updated 12/19/2022. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

<sup>\*</sup> Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (<a href="https://www.cdc.gov/std/treatment/default.htm">https://www.cdc.gov/std/treatment/default.htm</a>) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

## **List of Preferred Languages:**

**Instructions:** Complete the *Preferred Language* section if the patient's preferred language is <u>not</u> English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

## **Language Identity Codes:**

(L01) Amharic	(L18) Karen	(L35) Sign languages
(L02) Arabic	(L19) Khmer/Cambodian	(L36) Somali
(LO3) Balochi/Baluchi	(L20) Kinyarwanda	(L37) Spanish/Castilian
(LO4) Burmese	(L21) Korean	(L38) Swahili/Kiswahili
(L05) Cantonese	(L22) Kosraean	(L39) Tagalog
(L06) Chinese (unspecified)	(L23) Lao	(L40) Tamil
(L07) Chamorro	(L24) Mandarin	(L41) Telugu
(L08) Chuukese	(L25) Marshallese	(L42) Thai
(L09) Dari	(L26) Mixteco	(L43) Tigrinya
(L10) Farsi/Persian	(L27) Nepali	(L44) Ukrainian
(L11) Fijian	(L28) Oromo	(L45) Urdu
(L12) Filipino/Pilipino	(L29) Panjabi/Punjabi	(L46) Vietnamese
(L13) French	(L30) Pashto	(L77) Other language
(L14) German	(L31) Portuguese	(L88) Patient declined to respond
(L15) Hindi	(L32) Romanian/Rumanian	(L99) Unknown

(L33) Russian

(L34) Samoan

# **Race Category and Identity Instructions:**

**Instructions:** The patient's race may either be reported in the general *Race Category* section or the *Extended Race* section on page 1 of the case report. When completing the *Extended Race* section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the *extended race* section. **Unknown** race can be documented within either one of the race sections.

## **Extended Race Codes:**

(L16) Hmong

(L17) Japanese

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	